Table of Contents

ANDRUS, R. E. & D. H. VITT. *Sphagnotheca Boreali-americana*: A new exsiccata of North American sphagna north of Mexico 64

BARRETT, P. E. & J. W. THOMSON. Lichens from a high Arctic coastal lowland, Devon Island, N.W.T. 160

BASILE, D. V. A comparison of some macronutrient media used to culture bryophytes 403

BOERNER, R. E. & R. T. T. FORMAN. Salt spray and coastal dune mosses 57

BOWERS, F. D. & R. E. MAGILL. A new species of *Cyclodictyon* (Musci: Hookeriaceae) from Mexico 469

——— & A. J. SHARP. Additional mosses from Mexico of geographical significance 218

BRASSARD, G. R. New and additional moss records from Newfoundland 363

———. Major range extension for *Hygrohypnum polare* 488

——— & J. A. HANCOCK. Mosses of Fraser Canyon, Northern Labrador 484

BRODO, I. M. Review 237

CARMER, M.-B. Corticolous lichens of riparian deciduous trees in the central front range of Colorado 44

CHEVALLIER, D. Le manganèse: Oligoélément indispensable pour la germination des spores de *Funaria hygrometrica* 194

CROSBY, M. R. Recent literature on mosses—100–103 ..... 100, 225, 375, 491

CULBERSON, W. L. Recent literature on lichens—91–92 ..... 228, 380

DELGADILLO M., C. Taxonomic revision of *Aloina*, *Aloinella* and *Crossidium* (Musci) 245

DESMOND, M. E. & E. L. HARTMAN. Polyploidy and enucleation in mature leaves of *Barbilophozia barbata* 343

DOYLE, W. T. Spores of *Sphaerocarpos donnellii* 80

EGAN, R. S. Review 386

ELLIS, E. A. Observations on the ultrastructure of the lichen *Chiodecton sanguineum* 471

ENGEL, J. J. Hepaticae and anthocerotae collected by Dr. Harold E. Moore, Jr. in New Caledonia, Seychelles, Mauritius and Réunion in 1972 361

FAHSELT, D. Gas-liquid chromatography of the lichen substance usnic acid 452

FRENCH, J. C. & D. J. PAOLILLO, JR. Effect of exogenously supplied growth regulators on intercalary meristematic activity and capsule expansion in *Funaria* 431

——— & ———. On the role of the calyptra in permitting expansion of capsules in the moss *Funaria* 438

FRYXELL, P. A. Review 109

FULFORD, M. Recent literature on hepatics—70–72 ..... 96, 223, 372

GOUGH, L. P. Cryptogam distributions on *Pseudotsuga menziesii* and *Abies lasiocarpa* in the Front Range, Boulder County, Colorado 124

GRIFFIN, D. III. Additions to the moss flora of Venezuela 212

HOE, W. J. Additional new and noteworthy records for Hawaiian mosses—3 359

HONG, W. S. Hepaticae of the Black Hills, South Dakota 211

Table of Contents

ANDRUS, R. E. & D. H. VITT. *Sphagnotheca Boreali-americana*: A new exsiccata of North American sphagna north of Mexico 64

BARRETT, P. E. & J. W. THOMSON. Lichens from a high Arctic coastal lowland, Devon Island, N.W.T. 160

BASILE, D. V. A comparison of some macronutrient media used to culture bryophytes 403

BOERNER, R. E. & R. T. T. FORMAN. Salt spray and coastal dune mosses 57

BOWERS, F. D. & R. E. MAGILL. A new species of *Cyclodictyon* (Musci: Hookeriaceae) from Mexico 469

——— & A. J. SHARP. Additional mosses from Mexico of geographical significance 218

BRASSARD, G. R. New and additional moss records from Newfoundland 363

———. Major range extension for *Hygrohypnum polare* 488

——— & J. A. HANCOCK. Mosses of Fraser Canyon, Northern Labrador 484

BRODO, I. M. Review 237

CARMER, M.-B. Corticolous lichens of riparian deciduous trees in the central front range of Colorado 44

CHEVALLIER, D. Le manganèse: Oligoélément indispensable pour la germination des spores de *Funaria hygrometrica* 194

CROSBY, M. R. Recent literature on mosses—100–103 ..... 100, 225, 375, 491

CULBERSON, W. L. Recent literature on lichens—91–92 ..... 228, 380

DELGADILLO M., C. Taxonomic revision of *Aloina*, *Aloinella* and *Crossidium* (Musci) 245

DESMOND, M. E. & E. L. HARTMAN. Polyploidy and enucleation in mature leaves of *Barbilophozia barbata* 343

DOYLE, W. T. Spores of *Sphaerocarpos donnellii* 80

EGAN, R. S. Review 386

ELLIS, E. A. Observations on the ultrastructure of the lichen *Chiodecton sanguineum* 471

ENGEL, J. J. Hepaticae and anthocerotae collected by Dr. Harold E. Moore, Jr. in New Caledonia, Seychelles, Mauritius and Réunion in 1972 361

FAHSELT, D. Gas-liquid chromatography of the lichen substance usnic acid 452

FRENCH, J. C. & D. J. PAOLILLO, JR. Effect of exogenously supplied growth regulators on intercalary meristematic activity and capsule expansion in *Funaria* 431

——— & ———. On the role of the calyptra in permitting expansion of capsules in the moss *Funaria* 438

FRYXELL, P. A. Review 109

FULFORD, M. Recent literature on hepatics—70–72 ..... 96, 223, 372

GOUGH, L. P. Cryptogam distributions on *Pseudotsuga menziesii* and *Abies lasiocarpa* in the Front Range, Boulder County, Colorado 124

GRIFFIN, D. III. Additions to the moss flora of Venezuela 212

HOE, W. J. Additional new and noteworthy records for Hawaiian mosses—3 359

HONG, W. S. Hepaticae of the Black Hills, South Dakota 211

———. Leafy hepaticae of Montana and phytogeographic relationships to neighboring states and provinces 304
Ireland, R. R. *Homalothecium sericeum*, a neglected moss in North America 87
Johnsen, A. B. Additions to the hepatic flora of Arizona 200
Jørgensen, P. M. Further notes on Asian *Alectoria* 77
Kelly, B. B. & V. E. Becker. Effects of light intensity and temperature on nitrogen fixation by *Lobaria pulmonaria, Sticta weigelii, Leptogium cyanescens* and *Collema subfurvum* 350
Koch, R. G. Notes on Nebraska mosses II 85
Krog, H. & T. D. V. Swinscow. Variation in vegetative propagules in the African lichen *Parmelia lophogena* 67
Lipkin, Y. & V. W. Proctor. Notes on the subgenus *Trabutiella* of the aquatic liverwort *Riella* (Riellaceae, Sphaerocarpales) 25
Maass, W. S. G. Lichen substances VII. Identification of orsellinate derivatives from *Lobaria linita* 178
———. Lichen substances VIII. Phenolic constituents of *Pseudocyphellaria quercifolia* 183
McNeilus, V. E. & A. J. Sharp. *Preissia quadrata* in Tennessee 222
Manuel, M. G. Review of the genus *Penzigiella* (Bryopsida: Pterobryaceae) 423
———. Contribution to the bryoflora of Pennsylvania 458
Miller, N. G. Review 387
——— & K. M. Parks. Review 108
Mueller, D. M. J. Editorial 110
———. Review 239
Nash, T. H. III. Index to volume 78 500
——— & A. B. Johnsen. Catalog of the lichens of Arizona 7
O'Flaherty, L. M., J. D. Ives & A. R. Ozimek. *Sphagnum fimbriatum* new to Illinois 455
Pike, L. H., W. C. Denison, D. M. Tracy, M. A. Sherwood and F. M. Rhoades. Floristic survey of epiphytic lichens and bryophytes growing on old-growth conifers in Western Oregon 389
Pursell, R. A. Distributional records of rare or infrequently collected species of mosses in Pennsylvania II 93
——— & W. J. Hoe. *Fissidens nothotaxofolius* (Bryopsida: Fissidentaceae). A new species from the Hawaiian Islands 476
Ramaut, J. L. & Mme M. Corvisier. Les acides lichéniques chez *Stereocaulon virgatum* 365
Rao, K. R., C. S. P. Naidu & N. M. Kondaiah. Chemical constituents of *Riccia plana* 216
Redon, J. Review 239
Reese, W. D. *Calymperes rubiginosum* (Mitt.) Reese, comb. nov. 92
Sayre, G. Illustrations of the lost hepatics of the Wilkes Expedition 204
Scheirer, D. C. Anatomical studies in the Polytrichaceae II. Histochemical observations on thickened lateral walls of hydroids of *Dendroligotrichum* 113
Schroeder, N. E., G. J. Schroeder & D. E. Anderegg. Catalog of the lichens of Idaho 32
Showman, R. E. Lichens as indicators of air quality around a coal-fired power generating plant 1

SIGAL, L. L. & D. R. TOREN. A new *Haematomma* from California 70
——— & ———. New distribution records of lichens in California II 467
SMITH, D. K. Three interesting and unusual mosses discovered in East Tennessee 370
SMITH, G. L. Neotropical Polytrichaceae I, II 201
———. Neotropical Polytrichaceae III 480
SNIDER, J. A. Three new species of *Archidium* from Africa 152
SPESSARD, L. L. Five mosses new to Illinois 86
STEERE, W. C. Kjeld Axel Holmen, 1921–1974 240
———. Occurrence of *Entodon cladorrhizans* in Alaska, with notes on geographical distribution of *E. concinnus* 334
———. A new exsiccata from the northernmost parts of North America: Steere and Holmen, *Bryophyta Arctica Exsiccata*, Fascicle 1, Numbers 1–50 356
———. *Lophozia incisa*: A curious reaction to desiccation 368
——— & Z. IWATSUKI. Validation of Pseudoditrichaceae Steere et Iwatsuki fam. nov. 489
——— & G. L. SMITH. The sporophyte of *Oligotrichum falcatum* 447
TAKAO, A. Plasmolysis of protonema in *Bartramia pomiformis* 146
THOMAS, R. J. Sesquiterpenoids in the sporophyte of *Lophocolea heterophylla* 75
THOMSON, J. W. Review 238
TOWNSEND, C. C. *Pseudoscleropodium purum* on two more tropical islands 73
TUCKER, S. C. & D. T. KOWALSKI. New state records of lichens from Northern California 366
TÜRK, R. & V. WIRTH. Über die SO_2-Empfindlichkeit einiger Moose 187
VITT, D. H. & R. E. ANDRUS. *Sphagnum aongstroemii* in North America 463
——— & C. D. HAMILTON. Taxonomic status of *Tomenthypnum falcifolium* 168
WATSON, M. A. Annual periodicity of incremental growth in the moss *Polytrichum commune* 414
WEBER, W. A. Two new species of *Lecanora*, section *Petrasterion*, with a key to North American species 206
WEIN, R. W. & J. E. SPEER. Lichen biomass in acadian and boreal forests of Cape Breton Island, Nova Scotia 328

Index to Volume 78 of The Bryologist[1,2]

Abietinella, 337; abietina, 337, 357, 486
Acarospora americana, 8; badiofusca, 8; carnegei, 8; cervina var. cinereoalba, 8; chlorophana, 8, 32; chrysops, 8; cineracea, 8; coloradiana, 8; epilutescens, 8; flava, 8; fuscata, 8; geogena, 8; geophila, 8; glaucocarpa, 162; nigromarginata, 8; peltastica, 8; peltata, 8; radicata, 8; schleicheri, 8, 33; smaragdula, 8; strigata, 8; tucsonensis, 8; washingtonensis, 8; xanthophana, 8
Acrochila caledonica, 361
Acroporium pungens, 218
Actinodontium standleyi, 218
Actinogyra polyrrhiza, 32
Africa, three new species Archidium from, 152*
Agyrophora lyngei, 162, 165
Air pollution: lichens as indicators of air quality around a coal-fired power plant, 1; response of mosses to sulfur dioxide fumigations, 187
Alaska, Entodon from, 334
Alectoria sect. Divaricatae, 77; subg. Bryopogon, 78; subg. Teretiuscula, 78; bicolor, 77, 79; capillaris, 33; chalybeiformis, 8, 33, 163, 165; **cornicularioides** nom. nov. 77–8*; dahlii, 79; fremontii, 8, 33, 393; glabra, 33, 130–32, 135, 141, 143, 393, 400; implexa, 33; imshaugii, 392, 397, 401; indonesica, 77, 79; jubata, 8, 33; lata, 392, 397; minuscula, 8, 163, 165; nidulifera, 8, 33; nigricans, 163; nitidula, 79, 163; ochroleuca, 163, 165; oregana, 33, 393, 400–1; pubescens, 33, 77, 163; sarmentosa, 33, 393, 398, 400–1; subdivergens, 163; tenerrina, 33; tenuis, 163
Aloina, generic characters, 245–250; 279–83, 285–91, 295–6, 297, 301; aloides, 247–50, 255, 257, 292; aloides var. aloides, 247, 249–250, 255–8*, 284, 290, 292–3; aloides var. ambigua, 247, 249–250, 254, 256*–8, 284, 290, 292–3, 301; ambigua, *257*; anthropophila, *261*; apiculata, 248–250, 252*, 254–5, 284, 290; bifrons, 248, 250–3*, 284, 290, 293, 295, 301; brevirostris, 248–9, 251, 256*, 258–9, 262, 284, 286*–7, 289–91, 295–6, 301; brevirostris var. breidleri, *258–9*; brevirostris var. elongata, *259*; brevirostris var. rotundifolia, *259*; brevirostris var. typica, *259*; calceolifolia, 260, *261*, 262; catillum, 246, 248–9, 251, 256*, 260, 262, 284, 287, 290, 299; collina, *261*; **cornifolia** nom. nov., 245, 249, 251, 263*, 265, 284, 290; cucullatifolia, *268*; cucullifera, 268–9; ericaefolia, *257*; ericaefolia var. macrophylla, *257*; hamulus, 245, 248, 250, 252*, 254–5, 284, 287, 290, 296; longirostris, *254*, *261*; macrorhyncha, 259, *261*; obliquifolia, *264*; pilifera, *251*; potaninii, *261*, 265; recurvipatula, 246, 262, 279; rigida, 245, 248–9, 256, 259–262, 265, 285, 288, 291, 299–300; rigida ssp. aloides, 255; rigida var. ambigua, *257*; rigida var. longirostris, *262*; rigida var. mucronata, 245, 251, *262*; rigida var. mucronulata, 262*–5, 284, 290, 300–1; rigida var. obliquifolia, 245, 251, 263*–5, 284, 290; rigida var. obtusa, *261–2*; rigida var. pilifera, *251*; rigida var. rigida, 251, 256*, 260, 262, 264, 284, 290, 299–301; roseae, 249–250, 252*–3, 284, 287, 290, 299; sedifolia, 246, 262, 279; stellata, *261*; stellata var. obtusa, 261; stellata var. pilifera, *251*; sullivaniana, 251
Aloinella sp., 214, 279–81, 285–91, 301, generic characters, 245–250; andina, 248, 266–7, 287; apiculata, *254*; boliviana, 248, 266, 269, 272*; catenula, 245, 248, 265–8, 285–8*, 290, 301; catenulata, 214; cucullatifolia, 246, 266, 268–9, 272*; cucullifera, 248, 266, 268; galeata, 246, 279; hamulus, *254*
Amblystegium serpens, 486
Amphidium cyathicarpum 218; lapponicum, 286
Anacolia intertexta, 218
Anaptychia comosa, 8; diademata, 9; hypoleuca, 9; leucomelaena, 9; neoleucome-

[1] Compiled by T. H. Nash III, Department of Botany and Microbiology, Arizona State University, Tempe, Arizona 85281.

[2] New scientific names are printed in **boldface** type; illustrations are indicated by an asterisk; synonyms, by italicized page numbers.

laena, 9; obscurata, 9; pseudospeciosa, 9; pseudospeciosa var. tremulans, 9; speciosa, 9; squamulosa, 9; tropica, 9
Anastrepta, 309
Anastrophyllum, 309, 314; assimile, 315; donianum, 315; hellerianum, 315, 325; michauxii, 315, 325; minutum, 211, 315, 325
Anatomical studies, of Polytrichaceae, 113
ANDEREGG, D. E. *See* SCHROEDER, N. E., G. L. SCHROEDER and D. E. ANDEREGG
Andreaea crassinervia, 363; rupestris, 485
ANDRUS, R. E. and D. H. VITT: Sphagnotheca Boreali-americana: a new exiccata of North American sphagna north of Mexico, 64; *See* VITT, D. H. and R. E. ANDRUS
Anomobryum semiovatum, 214, 218–9
Anomodon minor, 219, 460; minor var. inaequifolius, 218–9; rostratus, 363; viticulosus, 218–9
Anthelia, 322; julacea, 322, 325; juratzkana, 322, 325
Anthoceros crispulus, 458–9
Anthocerotae, from New Caledonia, Seychelles, Mauritius and Réunion, 361
Antitrichia californica, 394, 399; curtipendula, 394, 399
Aongstroemia cucullata, 268; julacea, 221
Aphanorrhegma serratum, 85, 370, 460
Aptychella proligera, 218–9
Archidium **acanthophyllum** nom. nov., 152–4*, 157*; **andersonianum** nom. nov., 152, 156*–9; capense, 158; falcatulum, 158; **muellerianum** nom. nov., 152, 154–5*, 157*, 159; rehmannii, *154*; rothii, 154
Arctic, lichens of Devon Island, 160
Arizona: lichen catalog of, 7; additions to the hepatic flora, 200; **Lecanora christoi** from, 206
Arthonia, 124; mediella, 129–30, 133; radiata, 9
Aspicilia, 209; quartzitica, 9
Aspiromitus fuscus, 361
Atrichum altecristatum, 85, 363; angustatum, 85, 456; crispum, 460; haussknechtii, 363; oerstedianum, 363; undulatum, 85, 363; undulatum var. selwyni, 243
Aulacomnium acuminatum, 358; androgynum, 391–2; palustre, 57–62, 176, 357–8, 486; turgidum, 357, 484, 486

Bacidia sp., 47, 48, 51; abbrerians, 51; bagliettoana, 161; chlorococca, 33; fuscorubella, 33; herrei, 393; idahoensis, 33; naegelii, 393, 399; rubricosa, 33; sphaeroides, 33; vermifera, 48, 50, 51
Baeomyces carneus, 162; rufus, 33, 366
Baldwiniella kealeensis, 359
Barbilophozia, 348; barbata, 343, 345, 347*
Barbula, 246, sp. 391–2; acuta, 85; aloides, *255*; ambigua, *257*; anthropophila, *261*; breidleri, *258*; brevirostris, *258*; calceolifolia, *261*; catillum, *260*; chloronotos, 274, *276*; convoluta, 93; cucullatifolia, 246, *268*; cucullifera, *269*; duthieana, *261*, 288; fallax, 6; galeata, 246; grisea, *277*; hamulus, *254*; macrorhyncha, *261–2*; membranifolia, 274, *276*; membranifolia var. grisea, *277*; membranifolia var. laevipila, 278; membranifolia var. pottioidea, *277*; michiganensis, 218–9; obliquifolia, *264*; reflexa, 218–9; rigida, *260*; rigida var. bifrons, *251*; rigida var. brevirostris, *258*; rigida var. pilifera, *251*; rigida var. mucronata, *262*; rigida var. mucronulata, *262*; rigidum, 260; squamifera, 274, *276*; squamifera ssp. grisea, *277*; sullivaniana, *251*; unguiculata, 359
BARRETT, P. E. and J. W. THOMSON: Lichens from a high arctic coastal lowland, Devon Island, N.W.T., 160
Bartramia angustifolia, 214; glaucula, 221; halleriana, 218–9; ithyphylla, 486; microstoma, 214, 221; pomiformis, 146–151, 456
BASILE, D. V.: A comparison of some macronutrient media used to culture bryophytes, 403
Bazzania, 308; bernieri, 361; convexa, 361; decrescens, 361; deplanchei, 361; stephanii, 361; tricrenata, 458–9; trilobata, 459
BECKER, V. E. *See* KELLY, B. B. and V. E. BECKER
Benzyladenine, effect on Funaria, 432
Biatorella microhaema, *55*; moriformis, 9
Biatoria crenata, 9; icterica, 9; rufonigra, 9
Bilimbia rubicosa, 33; sphaeroides, 33
Biomass, of lichens from Nova Scotia, 328
Blastenia atrosanguinea, 9
Blepharostoma, 307; arachnoideum, 307, 326; trichophyllum, 211, 305–7; trichophyllum subsp. brevirete, 307, 325
Blindia acuta, 484–5
BOERNER, R. E. and R. T. T. FORMAN: Salt spray and coastal dune mosses, 57
BOWERS, F. D. and A. J. SHARP: Additional mosses from Mexico of geographical significance, 218; and R. E. MAGILL: A new species of Cyclodictyon (Musci: Hookeriaceae) from Mexico, 469
Brachymenium consimile, 218–9
Brachytheciaeae, 87
Brachythecium albicans, 62; campestre, 86, 456–7; curtum, 484, 486; oxycladon, 89; plumosum, 486; rivulare, 486; salebrosum, 370; velutinum, 94
BRASSARD, G. R.: New and additional moss

records from Newfoundland, 363; Major range extension for Hygrohypnum polare, 488; and J. A. HANCOCK: Mosses of Fraser Canyon, Northern Labrador, 484
Breutelia chrysea, 214; subarcuata, 214
BRODO, I. M. (review) D. H. S. Richardson, The Vanishing Lichens, 237
Bryoandersonia illecebra, 461
Bryoerythrophyllum recurvirostre, 93, 213; recurvirostrum, 85
Bryopsida, 423, 477
Bryum, 114; algovicum, 485; baldwinii, 360; capillare, 370, 393; longisetum, 485; pendulum, 62; procerum, 214; pseudocapillare, 359; rigidum, *260*; salinum, 485
Buellia atrata, 163; badia, 9, 33; betulina, 33; blumeri, 33; disciformis, 9, 33; glaziouana, 9; griseovirens, 33; lepidastra, 9; nigra, 33; novomexicana, 9; papillata, 163; parasema, 9, 33; penichra, 33, 393; pullata, 9; punctata, 9, 33, 48, 51, 130–2, 135, 140–1; retrovertens, 9; semitensis, 9; spuria, 9; tucsonensis, 9; turgescens, 9; zahlbruckneri, 9, 33
Buelliella inquilina, 33
Buxbaumia indusiata, 363; minakatae, 363

Calicium abietinum, 33; hyperellum, 33; pusillum, 392, 397; trabinellum, 9; viride, 33, 392, 397
California, **Lecanora mellea** from, 208; new lichens from, 366, 467
Callicladium haldanianum, 456–7
Calliergon sarmentosum, 486; stramineum, 486; trifarium, 215
Calliergonella cupidata, 337
Caloplaca sp., 48–51; amabilis, 9; arizonica, 9; atroalba, 9; atrosanguinea, 9; aurantiaca, 9, 33; cerina, 9, 33, 48, 51, 52, 54; chrysophthalma, 9; cinnabarina, 9; cinnamomea, 163; decipiens, 9; discolor, 9; durietzii, 9; elegans, 10, 33; epithallina, 10; ferrugniea, 10, 33; festiva, 10; flavovirescens, 10; fraudans, 10; fulgens, 10; gilva, 10; herbidella, 130, 133, 141; holocarpa, 10, 47, 48, 51, 163; jungermanniae, 10, 33, 163; lobulata, 10; microphyllina, 10; modesta, 10; murorum, 10, 33; oregona, 33; pinicola, 10; pyracea, 10; saxicola, 10; sideritis, 10; sorocarpa, 47, 48, 50, 51; spaldingii, 10; squamosa, 10; stillicidiorum, 163; tetraspora, 163; tiroliensis, 163; trachyphylla, 10; verrucosa, 10; wrightii, 10
Calymperes **rubiginosum** comb. nov., 92; rufum, *92*
Calypogeia, 308; fissa, 309, 458–9; integristipula, 309, 325; meylanii, 325; muelleriana, 305, 309, 325; neesiana, 309; sphagnicola, 309, 325; suecica, 309, 326; sullivantii, 460–1; trichomanis, 308–9, 325
Calyptra, role of in Funaria, 438
Camptothecium, 168; lutescens, 87, 89; nitens var. falcifolium, 172; nitens var. falcatum, 172; sericeum, *89*
Campylium chrysophyllum, 370; chrysophyllum var. brevifolium, 94; hispidulum, 215; stellatum, 176
Campylopus annotinus, 213; argyrocaulon, 213; savannarum, 213
Canada, lichens of Devon Island, 160; hepatics of, 304
Candelaria concolor, 10, 33, 48, 52, 130, 133, 141; fibrosa, 10
Candelariella sp., 48, 52; antennaria, 34; arctica, 162; aurella, 10, 47, 48, 52, 54, 129–30, 132, 141, 165; cerinella, 34; deflexa, 10; medians, 10; rosulans, 10, 34; submexicana, 10; terrigena, 10, 162; vitellina, 10, 34, 47–8, 52, 130–1, 133, 140–1; xanthostigma, 10
Capsules, expansion of in Funaria, 431, 438
CARMER, M.-B.: Corticolous lichens of riparian deciduous trees in the central front range of Colorado, 44
Catagonium politum, 218–9
Catillaria atropurpurea, 34; glauconigrans, 10; lenticularis, 10
Catoscopium nigritum, 357
Cephalozia, 322; ambigua, 322, 325; bicuspidata, 322, 325, 459; catenulata, 322, 459; connivens, 322, 325; lacinulata, 322, 326; lammersiana, 322; leucantha, 322, 325; loitlesbergeri, 322–3, 326; lunulifolia, 305, 322–3, 325; macounii, 322–3, 326; media, 325; pleniceps, 306, 322–3, 325
Cephaloziella alpina, 324; arctica, 324; brinkmani, 324; dentata, 324; divaricata, 324–5; hampeana, 324, 326; phyllacantha, 324; rubella, 211, 305, 324, 326, 393; subdentata, 324–5; turneri, 324
Cephaloziopsis, 324
Ceratodon purpureus, 57–60, 62–3, 359, 485
Cetraria atlantica, 34; californica, 34; canadensis, 34; chlorophylla, 34, 393; coralligera, 10; cucullata, 162, 165–6; delisei, 162, 165–6; ericetorum, 162; fendleri, 10, 130, 132, 141; fendleri fo. coralligera, 10; glauca, 34; idahoensis, 34, 366, 392; islandica, 10, 162, 166; juniperina, 34; merrillii, 34; nivalis, 162, 165–6; orbata, 34, 393; pallidula, 34, 393; pinastri, 10, 34, 130, 132, 135, 140–1, 143; platyphylla, 34, 393; subalpina, 34; tristis, 34; weberi, 10

Chaenotheca brunneola, 34; chrysocephala, 10, 34; ferruginea, 392, 397; trichialis, 34
Chandonanthus setiformis, 357
Chemistry, of Lophocolea heterophylla, 75; of Lobaria linita, 178; of Pseudocyphellaria quercifolia, 183; of Stereocaulon virgatum, 365
Chenothecopsis, 392
CHEVALLIER, D.: Le Manganèse: oligoélément indispensable pour la Germination des Spores de Funaria hygrometrica, 194
Chiloscyphus, 319–20; sp., 361; crispata, 205; fragilis, 200; pallescens, 200, 320, 325; pallescens var. fragilis, 320, 325; polyanthus, 200, 211, 305–6, 320, 325; polyanthus var. rivularis, 211, 320–1, 325; porrectus, 205; rivularis, 200
China, Alectoria cornicularioides from, 77
Chiodecton sanguineum, 471–2, 473*, 474*, 475
Chloronotus, 246
Chromatography, gas-liquid of usnic acid, 452
Cinclidium arcticum, 357; subrotundum, 357
Cladina alpestris, 330; mitis, 34, 330
Cladonia sp., 130, 132, 141; amaurocraea, 162; bacillaris, 10, 34, 124, 129, 141; balfourii, 11; bellidiflora, 162; borbonica fo. cylindrica, 34; capitata, 11; cariosa, 11, 34; carneola, 34; cenotea, 11, 34, 366; chorophaea, 11, 34, 141; chorophaea fo. carpophora, 34; chorophaea fo. prolifera, 34; chorophaea fo. simplex, 34; coccifera, 34, 162; coniocraea, 11, 34, 130, 133, 141; coniocraea fo. ceratodes, 11, 34; coniocraea fo. truncata, 11, 34; cornuta, 11, 34, 162; cornutoradiata, 34; crispata, 34; cristatella, 472; deformis, 11; digitata, 34; ecmocyma, 34; fimbriata, 11, 34, 141, 393; fimbriata var. chlorophaeoides, 34; fimbriata var. macra, 34; fimbriata var. simplex, 34; formosana, 34; furcata, 34; glauca, 35; gonecha, 35; gracilis, 35, 162, 165; lepidota, 162; macilenta, 35, 392, 397; major, 35; major fo. denticulata, 35; mitis, 162, 165; mitrula, 35; multiformis, 35; multiformis fo. finkii, 35; multiformis fo. simulata, 35; multiformis fo. subascypha, 35; nemoxyna, 35; ochrochlora, 35; phyllophora, 35; pityrea, 35; pleurota, 35; pocillum, 162, 166; polydactyla, 35; pyxidata, 11, 35, 141; scabriuscula, 35; squamosa, 35; subsquamosa, 392; subulata, 35; theiophila, 35; turgida, 35; verticillata, 35; yunnana, 35
Clapodium bolanderi, 394
Cladopodiella, 322
Climacium dendroides, 486
Coccocarpia cronia, 11
Collema ceraniscum, 161; coccophorum, 11; conglomeratum, 11; conglomeratum var. crassiusculum, 11; furfuraceum, 11, 47–8, 52, 366; polycarpon, 11; subfurvum, 11, 350–5; tenax, 11; texanum, 11; tunaeforme, 11, 366; undulatum, 11; undulatum var. granulosum, 161; verruciforme, 11
Cololejeunea biddlecomiae, 459
Colorado: lichens of riparian deciduous trees of the Central Front Range, 44; lichens on Pseudotsuga menziesii and Abies lasiocarpa in Boulder County, 124; **Lecanora christoi** from, 206
Coniocybe furfuracea, 392, 397
Conocephalum conicum, 211
Conostomum, 363; tetragonum, 363, 486
Cora pavonia, 472
Coriscium viride, 163
Cornicularia, 77; aculeata, 163, 367; divergens, 163; muricata, 163; normoerica, 35
CORVISIER, MME M. *See* RAMAUT, J. L. and MME M. CORVISIER
Crocynia lanuginosa, 11, 35; membranacea, 11
CROSBY, M. R.: Recent literature on mosses, 100, 100; 101, 225; 102, 375; 103, 491
Crossidium, generic characters, 245–50; 279–83, 285, 287–91, 294, 298, 301; aberrans, 245, 248, 269, 271–2*, 284, 290, 293–4; chloronotos, 274–6; chloronotos var. niveum, 276; crassinerve, 245, 248–9, 274–5, 279, 285, 290, 293, 297; crassinerve var. crassinerve, 270, 272*, 274, 276, 284; crassinerve var. laevipilum, 245, 270, 275–6, 284; criseum, 277; desertorum, 274–5; elatum, 278; erosum, 274–5; geheebii, 248–9, 270–2*, 284–5, 290, 294; laevipilum, 275; membranifolium, 276; peruvianum, 253; roseae, 253; rosei, 248–9, 269, 272*–4, 284–5, 290, 294; seriatum, 248–9, 270–2*, 284, 290, 294; spatulaefolium, 270–1; squamiferum, 247–9, 269, 274, 276, 279, 285; squamiferum ssp. griseum, 277; squamiferum ssp. squamiferum, 276, 284; squamiferum var. brevisetum, 278–9; squamiferum var. griseum, 277; squamiferum var. longipilum, 278–9; squamiferum var. pottioideum, 247–9, 270, 272*, 277–8, 284–5, 290, 293, 298; squamiferum var. pumilum, 278–9; squamiferum var. squamiferum, 247–8, 270, 272*, 276–7, 290, 293, 298; succulentum, 277
Ctenidium molluscum, 187–8, 190–2, 460
CULBERSON, W. L.: Recent literature on lichens, 91, 228; 92, 380
Culture, of mosses, 403
Cyclodictyon, 469; albicans, 470; richardsii, 469–70*

Cylindrothecium, 337–8
Cynodontium alpestre, 485; strumiferum, 485
Cyphelium inquinans, 35, 392, 397; lucidum, 11; tigillare, 11
Cyrtomnium hymenophyllum, 358

Dactylina arctica, 163, 165–6; ramulosa, 163, 166
Daltonia, 471; gracilis, 218–9; latolimbata, 215; peruviana, 215; pulvinata, 218–9; stenophylla, 218, 220; tenuifolia, 218, 220
Dawsonia, 113
Delgadillo, M. C.: Taxonomic revision of Aloina, Aloinella and Crossidium (Musci), 245
Dendroalsia abietina, 394
Denison, W. C. *See* Pike, L. H., et al.
Dentroligotrichum, 113–22; dendroides, 114–5*
Dermatocarpon acarosporoides, 11; aquaticum, 35; compactum, 11; fluviatile, 11; hepaticum, 35; intestiniforme, 11; lachneum, 11; miniatum, 11; miniatum var. papillosum, 35; moulinsii, 11; moulinsii var. subpapillosum, 35; peltatum, 11; plumbeum, 11; polyphyllum, 11; reticulatum, 35, 367; rufescens, 11; rupicola, 11; squamellum, 367–8; tuckermanii, 11
Dermea sp., 129, 131–2, 143
Desiccation, of Lophozia incisa, 368
Desmatodon, 281–3; convolutus, 281; griseus, 277–8; heimii, 363
Desmond, M. E. and E. L. Hartman: Polyploidy and enucleation in mature leaves of Barbilophozia barbata, 343
Development, effect of growth regulators in Funaria, 431
Dicranoweisia cirrata, 192, 391–2; crispula, 485
Dicranum spp., 484; angustum, 357; bonjeanii, 485; elongatum, 485; fragilifolium, 485; fuscescens, 391, 393, 397–8, 401, 485; groenlandicum, 485; howellii, 394; majus, 485; muehlenbeckii, 485; scoparium, 57, 343; tauricum, 393
Didymodon asperifolius, 489; recurvirostris, 93
Dimelaena oreina, 12; thysanota, 12
Dimerella diluta, 35
Diplophyllum albicans, 317; obtusifolium, 317; taxifolium, 317, 325, 393, 399
Diploschistes actinostomus, 35; actinostomus var. brunneus, 12; bryophilus, 12; scruposus, 12, 35
Distribution of: corticolous lichens of riparian trees in the Central Front Range of Colorado, 44; Parmelia lophogena*, 67; Haematomma californicum, 72; Pseudoscleropodium purum, 73; Asian Alectorias, 77; Nebraska mosses, 85; Illinois mosses, 86; Homalothecium sericeum, 87; Calymperes rubiginosum, 92; Pennsylvania mosses, 93; lichens on Pseudotsuga menziesii and Abies lasiocarpa in Boulder County, Colorado, 124; three species of Archidium, 152*; Tomenthypnum, 168; hepatics in Arizona, 200; Polytrichaceae, 201; North American Lecanoras in Sect. Petrasterion, 206; mosses from Venezuela, 212; mosses from Mexico, 218; hepatics of Montana and adjacent regions, 304; Entodon, 334; Aloina, Aloinella, and Crossidium, 245; mosses of Hawaii, 359; mosses from Newfoundland, 363; lichens in California, 366, 467; Penzigiella, 423; mosses in Labrador, 484; Oligotrichum falcatum, 447; the Bryopora of Pennsylvania, 459
Ditrichum flexicaule, 485; lineare, 93
Douinia, 317
Doyle, W. T.: Spores of Sphaerocarpos donnellii, 80
Drepanocladus exannulatus, 176, 215, 486; fluitans, 244; procerus, 176; revolvens, 176, 486; uncinatus, 173, 358, 486
Dryptodon patens, 363

Ecological studies: of corticolous lichens of riparian deciduous trees in the Central Front Range of Colorado, 44; salt spray and coastal dune mosses, 57; of lichens on Pseudotsuga menziesii and Abies lasiocarpa, Boulder County, Colorado, 124; of lichens and bryophytes of old-growth conifers of western Oregon, 389
Ecuador, Polytrichastrum from, 201
Editorial, 110
Egan, R. S. (review) M. E. Hale, Jr. The Biology of Lichens, 386
Electron micrographs: of Dendroligotrichum dendroides, 115–6; of Chiodecton sanguineum, 473–4 *See* SEM
Ellis, E. A.: Observations on the ultrastructure of the lichen Chiodecton sanguineum, 471
Encalypta spp., 484; brevicolla, 485; ciliata, 485; procera, 460; streptocarpa, 460
Endocarpiscum placodizans, 12
Endocarpon pusillum, 12; schaereri, 12
Engel, J. J.: Hepaticae and Anthocerotae collected by Dr. Harold E. Moore, Jr. in New Caledonia, Seychelles, Mauritius and Réunion in 1972, 361
Entodon, 337; brevisetus, 363; cladorrhizans, 334–5; concinnus, 334–8, 340, 357, 363; concinnus subsp. caliginosus, 337; con-

cinnus subsp. concinnus, 337; concinnus var. fauriei, 338; montagnei, 338; orthocarpus, 336, 338
Eriopus setigerus, 215
Eurhyndhium hians, 428; pulchellum, 215; vagans, 259
Evernia divaricata, 12; furfuracea, 12; furfuracea var. ceratea, 12; prunastri, 35; vulpina, 12, 35
Exsiccata: of North American Sphagna, 64; of bryophytes from northern North America, 356

Fabronia nietneri, 359
Fahselt, D.: Gas-liquid chromatography of the lichen substance usnic acid, 452
Fissidens adelphinus, 476, 478; bishopii, 360; bushii, 476, 478; cristatus, 363–4; crumii, 360; hawaiicus, 359; kilaueae, 359; nothotaxifolius, 476–9*; osmundoides, 485; taxifolius, 478
Fissidentaceae, 477
Floristics: of cryptogams on old-growth conifers in western Oregon, 389
Fontinalis dalecarlica, 486
Forman, R. T. T. *See* Boerner, R. E. and R. T. T. Forman
Fossombronia, 368–9; foveolata, 368–9
French, J. C. and D. J. Paolillo, Jr.: Effect of exogenously supplied growth regulators on intercalary meristematic activity and capsule expansion in Funaria, 432; on the role of the calyptra in permitting expansion of capsules in the moss Funaria, 440
Frullania, 337; bolanderi, 394; eboracensis, 459; nisquallensis, 205, 394, 399–400; riparia, 211; tamarisci, 337; tamarisci subsp. nisquallensis, 337
Fryxell, P. A.: (review) Memoria del II Congreso Venezolano de Botanica (celebrado en Mérida entre junio 26 y julio 1 de 1972)
Fulford, M.: Recent literature on hepatics, 70, 96; 71, 223; 72, 372
Fulgensia bracteata, 12, 163, 165–6; fulgens, 12
Funaria, 114, 194–8, 431, 435–6, 438, 444; americana, 370; hygrometrica, 194, 198, 359, 432, 439*; lindigii, 214

Geocalyx, 319, 321; graveolens, 321, 325
Germination, of Funaria hygrometrica, 194
Globulinella peruviana, 218, 220
Glossadelphus zollingeri var. filicaulis, 359
Gonohymenia, 472
Gough, L. P.: Cryptogam distribution on Pseudotsuga menziesii and Abies lasiocarpa in the Front Range, Boulder County, Colorado, 124
Graphina parilis, 367
Graphis striatula, 367
Griffin III, D.: Additions to the moss flora of Venezuela, 212
Grimmia agassizii, 484–5; alpestris, 364; alpicola, 485; donniana, 484–5; funalis, 484–5; ovalis, 485; pulvinata, 187–8, 192–3; tenerrima, 363–4; torquata, 364, 484–5
Growth: yearly of Polytrichum commune, 414; regulators in Funaria, 431
Gyalecta foveolaris, 161; peziza, 161
Gymnocolea, 309, 315; inflata, 315, 325
Gymnomitrion concinnatum, 317, 325; coralloides, 317; obtusum, 317, 326; pacificum, 317; varians, 317
Gymnostomum aeruginosum, 94; calcareum, 94; recurirostre, 94
Gyrophora decussata, 35; decussata var. darrowii, 35; dillenii, 35; erosa, 35; flocculosa, 35; hirsuta, 12; hyperborea, 36; mühlenbergii, 36; phaea, 36; proboscidea, 12, 36; rugifera, 12, 36; torrefacta, 36; vellea, 36

Haematomma **californicum** nom. nov., 70–2; lapponicum, 36, 70–1, 162; leiphaemum, 71; subpuniceum, 12
Hancock, J. A. *See* Brassard, G. R. and J. A. Hancock
Haplohymenium triste, 220
Harpanthus, 319, 321; flotovianus, 321, 325; scutatus, 321
Hartman, E. L. *See* Desmond, M. E. and E. L. Hartman
Hawaii, new mosses from, 359; Fissidens nothotaxifolius from, 476
Hepaticae, 459
Hepatics, notes on Riella, 25; additions to Arizona, 200; from Wilkes Expedition*, 204; of the Black Hills, South Dakota, 211; from Montana and neighboring regions, 304; from New Caledonia, Seychelles, Mauritius and Réunion, 361
Heppia bolanderi, 12; deserticola, 12; euploca, 12; lutosa, 12, 367–8; placodizans, 12; polyspora, 12; tortuosa, 12; virescens, 12
Herberta mascarenica, 361
Herpothallon sanguineum, 475
Herzogiella turfacea, 94
Heterodermia galactophylla, 12; hypoleuca, 12; leucomela, 12; obscurata, 12; pseudospeciosa, 12; squamulosa, 12; tremulans, 12
Hoe, W. J.: Additional new and noteworthy records for Hawaiian mosses—3, 359; *see* Pursell, R. A. and W. J. Hoe
Holosteum umbellatum, 293
Homalothecium, 168–9; fulgescens, 87, 394;

megaptilum, 168; nitens, 91; nitens var. falcifolium, 172; nuttallii, 90–1; **sericeum** comb. nov., 87–91*
HONG, W. S.: Hepaticae of the Black Hills, South Dakota, 211; leafy hepaticae of Montana and phytogeographic relationships to neighboring states and provinces, 304
Hookeria acutifolia, 220, 359
Hookeriaceae, 469
Hookeriopsis incurva, 218, 220
Hydrothyria venosa, 472
Hygrobiella, 322–3; laxifolia, 323, 325
Hygrohypnum alpestre, 486; dilatatum, 484, 486; eugyrium, 484, 486; ochraceum, 486; polare, 358, 488; range extension of, 488
Hylocomium brevirostre, 218, 220; splendens, 176, 394, 415, 484; splendens var. obtusifolium, 484, 486; splendens var. splendens, 486
Hymenostylium recurvirostrum, 94
Hyophila involuta, 94, 458, 460; totula, 94
Hypnaceae, 423
Hypnum circinale, 391, 393, 397–9; concinnum, 336; cordatum, *427*, *428*, 429; cupressiforme, 129, 131, 133, 135, 143; fertile, 486; lindbergii, 486; nitens, 168; orthocarpon, 336; polypterum, 215; procerrimum, 489; revolutom, 486; sericeum, 89; subimponens, 394
Hypogymnia atrofusca, 12; austerodes, 130, 133, 141; bitteri, 12; bitteriana, 130, 133, 141; duplicata, 36; enteromorpha, 36, 392, 398–400; imshaugii, 36, 392, 399–400; inactiva, 393; intestiniformis, 36; physodes, 12, 36, 162, 393; subobscura, 162, 165; tubulosa, 36, 393; vittata, 36

Icmadophila ericetorum, 36, 367
Idaho: lichen catalog of, 32; hepatics of, 304
Illinois, five mosses from, 86; Sphagnum fimbriatum from, 455
Indoleacetic acid, effect on Funaria, 432
Indonesia, Alectoria indonesica and A. dahlii from, 79
Ionapsis epulotica, 12, 162
IRELAND, R. R.: Homalothecium sericeum, a neglected moss in North America, 87
Isodrepanium lentulum, 220
Isopaches, 309
Isopterygerium distichaceum, 94; muellerianum, 94; turfaceum, 94
Isopterygiopsis muelleriana, 94
Isopterygium pulchellum, 486
Isothecium cristatum, 393; sericeum, 89, 391; stoloniferum, 394
IVES, J. D. *See* O'FLAHERTY, L. M. et al.

Jamesoniella, 315; autumnalis, 305, 315, 325
Japan, Alectoria nitidula from, 79
JOHNSEN, A. B.: Additions to the hepatic flora of Arizona, 200; *See* NASH III, T. H. and A. B. JOHNSEN
JØRGENSEN, P. M.: Further notes on Asian Alectoria, 77
Jungermannia allenii, 315; decurvifolia, 205; humilis, 205; lanceolata, 305–6, 315, 325; quadripartita, 205; quanperlita, 205; schraderi var. antarctica, 205

KELLY, B. B. and V. E. BECKER: Effects of light intensity and temperature on nitrogen-fixation by Lobaria pulmonaria, Sticta weigelii, Leptogium cyanescens and Collema subfurvum, 350
KOCH, R. G.: Notes on Nebraska mosses II, 85
Koerberia biformis, 12, 467–8
KOWALSKI, D. T. *See* TUCKER, S. C. and D. T. KOWALSKI
KROG, H. and T. D. V. SWINSCOW: Variation in vegetative propagules in the African lichen Parmelia lophogena, 67
Kurzia, 308

Labrador, mosses of, 484
Lasallia papulosa, 12; pensylvanica, 12; pustulata subsp. papulosa, 12
Lecania cyrtella, 52; cyrtellina, 47–50, 52; erysibe, 12
Lecanora, 50, 206, 209; alphoplaca, 13, 36, 209; alpina, 36; atra, 13, 36, 162; badia, 13, 162; baicalensis, 208; behringii, 162; bipruinosa, 13, 208–10; cadubriae, 129–31, 133, 136, 142; caesiocinerea, 13; caesiorubella, 36; caesiorubella subsp. glaucomodes, 13; caesiorubella subsp. saximontana, 13; calcarea, 13; candida, 162; carpinea, 13, 36, 130, 133, 142; cascadensis, 209; castanea, 162; cenisia, 13; chlarona, 13, 36; chlarotera, 13, 47–8, 52, 129–31, 133, 142; chloropolia, 13, 47–8, 52, 133, 135–6, 140, 142; **chrystoi** nom. nov., 206–7*, 210; chrysoleuca, 13, 36; cinerea, 13, 36; cinereorufescens, 36; circinata, 13; coilocarpa, 36; contorta, 13; diffracta, 13; dispersa, 13, 162; distans, 36; epibryon, 162, 165–6; epulotica, 13; expallens, 124, 129–31, 133, 135, 142; frustulosa, 13, 36, 162; garovaglii, 13, 209; hageni, 36, 47–8, 52–6; hypoptoides, 124, 129–30, 132, 142; insignis, 36; intricata, 162; melaena, 13; melanaspis, 13; melanophthalma, 13, 36, 162; **mellae** nom. nov., 207*–10; muralis, 13, 36, 206, 208–10; mutabilis, 13; nevadensis, 209; nigromarginata, 208–10; nordenskioldii, 162; novomexicana, 208–

10; oregana, 36; pacifica, 36; pallida, 13, 36; parisiensis, 13; peltata, 36; phaedrophthalma, 208; phaeobola, 392, 397; piniperda, 13, 36, 124, 129–30, 132, 142; polytropa, 13, 36, 162; proserpens, 162; radiosa, 13, 209; rubina, 13, 36; rugosa, 36; rugosella, 13, 36; rupicola, 13, 36, 162; subfusca, 13, 36; subfusca var. variolosa, 13; subfuscata, 37; subintricata, 48, 50, 52; subisabellina, 209; subrugosa, 37, 162; supertegens, 467–9; symmicta, 13, 37; thamnoplaca, 13; thomsonii, 208–10; urceolaria, 13, 37, 162; varia, 13, 37, 128–32, 135–6, 139–40, 142–3; verrucosa, 37

Lecidea, 53; anthracophila, 13; armeniaca, 14, 161; assimilata, 161; atrata, 161, atrobrunnea, 14, 37; auriculata, 37, 161; berengeriana, 14, 394; brandegei, 14; carnulenta, 37; carpathica, 14; cascadensis, 37; cinnabarina, 37, 393, 399; columbiana, 37; crassipes, 161; crenata, 14; crustulata, 161; cyanea, 37; decipiens, 14; elaeochroma, 14; elata, 14; elegantior, 14; enteroleuca, 37; erratica, 393, 399; euphorea, 14, 37, 53; friesii, 14, 37, 392, 397; fuscescens, 37; fuscoatra, 14, 37; glaucophaea, 161; globifera, 14, 37; glomerulosa, 14, 37, 130–2, 136, 142; granulosa, 14; icterica, 14; lapicida, 37, 161; latypiza, 14; lucida, 37; lulensis, 162; lurida, 14; lyngei, 14; macrocarpa, 14, 37, 162; melinodes, 162, 165; micacea, 162; novomexicana, 14, 37; ochrophora, 55; quadricolor, 14; paddensis, 467–8; pantherina, 162; parasema, 14, 37; platycarpa, 14; plebeja, 124, 129–30, 133, 142; purissima, 162; ramulosa, 162, 166; rubiformis, 14, 37, 162; rufonigra, 14; rugosella, 37; scalaris, 14, 37, 392, 397; speirea, 37, 162; stigmatea, 14; subaglaea, 14; subfusca, 37; symmicta, 14, 37; tesselina, 14; tessellata, 14, 37, 162; tornoensis, 14, 37, 392, 399; turgidula, 14; vernalis, 14, 37, 162; viridans, 14; viridescens, 14, 37; vorticosa, 14, 162; vulgata, 14

Lecidella carpathaca, 53; elaecochroma, 14; **glomerulosa** comb. nov., 44, 48, 50, 53; stigmatea, 14, 48, 53, 162, 165; wulfenii, 162

Leiciophysma finmarkicum, 161

Lejeunea sp., 361; **denudata** comb. nov., 361

Lepidozia, 308; filamentosa, 308; reptans, 211, 305, 308, 325; sandvicensis, 308

Lepraria sp., 48–9, 53, 130, 132, 142; chlorina, 14; membranacea, 14, 37, 392, 397; neglecta, 37, 163

Leprocaulon albicans, 14

Leptobryum pyriforme, 214, 485

Leptodontium, 247; brachyphyllum, 213; capituligerum, 213; erythroneuron, 213; longicaule var. longicaule, 213; luteum, 213; stellaticuspis, 213; subgracile, 213; wallisii, 213

Leptogium sp., 47–8, 53; apalachense, 14; arizonicum, 15; arsenei, 15; austroamericanum, 15; burgessii, 15; californicum, 37; chloromelum, 15; cyanescens, 14, 350–5; denticulatum, 15; furfuraceum, 15; hildenbrandii, 15; hypotrachynum, 15; inflexum, 15; juniperinum, 15; lacerum var. pulvinatum, 37; lichenoides, 15, 37; millegranum, 15; minutissimum, 161; palmatum, 37, 394; rugosum, 15; saturninum, 15; sinuatum, 394; tenuissimum, 15

Leptorhaphis epidermidis, 15

Lepyrodon tomentosus, 218, 220

Leskea obscura, 461; sericea, 87, 89; sericea var. striatula, 89

Letharia columbiana, 37, 392, 397; vulpina, 15, 37

Leucobryum glaucum, 456; seemannii, 360

Leucodon brachypus var. andrewsianus, 364

Leucolepis menziesii, 394

Lichen substances, orsellinate derivatives from Lobaria linata, 178; phenolic constituents of Pseudocyphellaria quercifolia, 183; chromatography of, 452

Lichens, as air quality indicators, 1; of Arizona, 7; of Idaho, 32; of deciduous trees in the Front Range of Colorado, 44; of Devon Island, 160; biomass in Nova Scotia of, 328; of California, 366; of old-growth conifers of western Oregon, 389; of California, 467

Lichina confinis, 350, 354

Lichinella americana, 15; stipatula, 15

Light intensity, effects on N-fixation in lichens, 350

LIPKIN, Y. and V. W. PROCTOR: Notes on the subgenus Trabutiella of the aquatic liverwort Riella (Riellaceae, Sphaerocarpales), 25

Lobaria dissecta, 185; hallii, 37; linita, 178–81; oregana, 37, 350, 391–2, 398*; pulmonaria, 37, 67, 330, 350–5, 393, 399; scrobiculata, 393, 399

Lopadium pezizoideum, 37, 162, 393

Lophocolea, 319–20; bidentata, 320; cuspidata, 320, 325; heterophylla, 75–6, 320, 325, 456, 459; longispica, 361; minor, 211, 320, 325

Lophozia, 309; subg. Barbilophozia, 309–10; subg. Leiocolea, 309–10; subg. Lophozia, 309, 312; subg. Massula, 309, 311; subg. Orthocaulis, 309; alpestris, 312, 325; ascendens, 211, 305, 213, 326; atlantica,

309, 325; attenuata, 309, 326; badensis, 310, 326; bantriensis, 305, 310, 325; barbata, 310, 325; binsteadii, 309; confertifolia, 312; excisa, 312, 325; floerkei, 309, 325; gillmani, 310–1, 325; grandiretis, 311; hatcheri, 306, 310, 325; heterocolpa, 305, 310–1, 325; incisa, 305, 311, 325, 368–9; kunzeana, 309, 326; longidens, 211, 305, 312–3, 326; lycopodioides, 306, 310, 325; marchica, 311; muelleri, 305–6, 310–1, 326; obtusa, 306, 311–2, 325; porphyroleuca, 305–6, 312–3, 325; quadriloba, 309, 325; rutheana, 310–1, 325; ventricosa, 211, 305, 312–3, 325; ventricosa var. longiflora, 312, 314, 325; wenzelii, 312, 314, 325
Lyellia, 453; aspersa, 451; crispa, 451

Maass, W. G. G.: Lichen substances VII. Identification of orsellinate derivatives from Lobaria linita, 178; VIII. Phenolic constituents of Pseudocyphellaria quercifolia, 183
Macrodiplophyllum, 317
Macromitrium emersulum, 360; shankii, *92*
Madotheca navicularis, 205
Magill, R. E. *See* Bowers, F. D. and R. E. Magill
Mala Kondaiah, N. *See* Rao, K. R., C. S. Prabhakar Naidu and N. Mala Kondaiah
Manganese, effects on germinations of Funaria hygrometrica, 194
Manuel, M. G.: Review of the genus Penzigiella (Bryopsida: Pterobryaceae), 423; Contribution to the Bryoflora of Pennsylvania, 458
Marchantia polymorpha, 211
Marsupella, 317; alpina, 317; brevissima, 317; commutata, 317; emarginata, 317, 325; sphacelata, 317, 326; stableri, 317
Massalongia microphylliza, 15
Mastigobryum fuegiana, 205
Mastigophora diclados, 361
Mattenccia struthiopteris, 146, 149–50
Mauritius, Anthocerotae and hepatics from, 361
McNeilus, V. E. and A. J. Sharp: Preissia quadrata in Tennessee, 222
Meesia triquetra, 357
Meiothecium boryanum, 218, 220
Meristem activity in Funaria, effects of growth regulators on, 432
Metaneckera menziesii, 394
Meteoriaceae, 425
Meteorium, 428; cordatum, *427*; hookeri, *427*, 429
Metzgeria decipiens, 361; furcata, 192
Mexico, Pogonatum from, 203; mosses of, 218; new species of Cyclodictyon, 469
Micarea, 53; populina, 48, 53–6; viridescens, 15, 37
Microthelia aterrima, 15; micula, 15
Mielichhoferia, 485; campylocarpa, 214; elongata, 485; mielichhoferi, 484–5
Miller, N. G.: (review) Academia Sinica (ed.) Iconographia Cormophytorum Sinicorum, 108; (review) P. D. Moore and D. J. Bellamy, Peatlands, 387
Mittenothamnium squarrosolum, 220
Mnium, 114; ciliare, 456; cuspidatum, 85, 456–7; orthorrynchum, 85; punctatum, 85; spinosum, 336; undulatum, 187–8, 190–2
Montana, hepatics of, 304
Morinia ehrenbergiana, 213
Morphological variation: of Parmelia lophogena, 67
Mosses, of Nebraska, 85; of Illinois, 86; of Pennsylvania, 93; response to sulfur dioxide, 187; of Venezuela, 212; of Mexico, 218; of Hawaii, 359; of Newfoundland, 363; of Tennessee, 370; of old-growth conifers of western Oregon, 389; culture techniques, 403; of northern Labrador, 484
Mueller, D. M. J. (review) S. R. Kashyap. Liverworts of the western Himalayas and the Panjab Plain, 239; Editorial, 110
Mycoblastus alpinus, 162; sanguinarius, 38, 162, 392, 398–9
Mycocalicium albonigrum, 38
Mylia, 315
Myrmecia, 351
Myurella julacea, 222, 357, 486
Myuroclada maximowiczii, 334

Nardia, 315; breidleri, 317, 325; compressa, 317, 326; geoscyphus, 317, 325, 459; geoscyphus var. insecta, 317; scalaris, 317, 325
Nash III, T. H. Index to Volume 78, 500–15; and A. B. Johnsen: Catalog of the lichens of Arizona, 7
Neckera cordata, *427*, *428*, 429; douglasii, 391, 394; novae-granadae, 214
Neckeraceae, 425; subfamily Thanmoideae, 425
Neodolichomitra, 423
Nephroma arcticum, 350; bellum, 15, 392; expallidum, 161; helveticum, 15, 38, 394; helveticum var. helveticum, 38; helveticum var. sipeanum, 38; laevigatum, 391, 394; parile, 15, 38, 367, 393; resupinatum, 15, 38, 391, 394
New Caledonia, Anthocerotae and hepatics from, 361
New Guinea, Alectoria indonesica and A.

dahlii from, 79; Entodon concinnus from, 334
Newfoundland, mosses of, 363
News and notes, 111, 242, 388, 498
Nigella, 293
Nitrogen-fixation, in Lobaria pulmonaria, Sticta weigelii, Leptogium cyanescens and Collema subfurvum, 350
Normandina pulchella, 15, 367, 393
North America, Sphagnum aongstroemii in, 463
Notothylas orbicularis, 460
Nova Scotia, lichen biomass of, 328
Nowellia curvifolia, 460

Ochrolechia androgyna, 15, 38, 162, 393; arborea, 38; frigida, 162; geminipara, 162; gonatodes, 162; inaequatula, 162, 165; oregonensis, 392, 397–9; pallescens, 15, 38, 393; parella, 338; upsaliensis, 38, 162, 165
Octoblepharum albidum, 360; pulvinatum, 220
O'Flaherty, L. M., J. D. Ives and A. R. Ozimek: Sphagnum fimbriatum new to Illinois, 455
Oligotrichum, 453; falcatum, 447, 448*, 449*–51; semilamellatum, 447, 451
Omphalodiscus decussatus, 15, 38; kraschenninikovii var. darrowii, 15, 38; virginis, 15, 38, 162
Omphalodium arizonicum, 15; hottentottum var. arizonicum, 15
Oncophorus virens, 485; wahlenbergii, 357, 485
Oreas martiana, 336
Oregon, cryptogams of old-growth conifers, 389
Oreoweisia bogotensis, 213; laxiretis, 213
Oropogon, 77
Orthothecium chryseum, 358
Orthotrichum aequatoreum, 214; chryseum, 169; chryseum var. falcifolium, 169; consimile, 394, 400; elongatum, 214; obtusifolium, 129, 131, 133, 144; pylaisii, 486; strangulatum, 370
Orsellinate derivatives, 178–81*, 183–86*
Osterwaldiella, 427
Ozimek, A. R. *See* O'Flaherty, L. M. et al.

Pallavicinia lyellii, 459–60
Paludella squarrosa, 357–8, 486
Pannaria hookeri, 161, 165; leucosticta, 467–8; microphylla, 15; pezizoides, 38; pityrea, 15, 367
Paolillo, Jr., D. J. *See* French, J. C. and D. J. Paolillo, Jr.
Parmelia ajoensis, 15; arseneana, 38; aspersa, 15, 38; aspidota, 15; aurulenta, 16; bolliana, 16; borreri, 16; caperata, 1–5, 16; centrifuga, 38, 162, 165; chiricahuensis, 16; chlorochroa, 16; conspersa, 16, 38; conspersa var. isidiata, 38; conspersa var. stenophylla, 38; crinita, 16, 367; cumberlandia, 16, 454; darrowii, 16; diacidula, 68*; disjuncta, 38, 162, 165; dissensa, 16; dolosa, 67–8; elegantula, 16, 38, 129–31, 133, 136, 140, 142; enteromorpha, 38; eurysaca, 16; exasperata, 16, 38; exasperatula, 38, 48, 53, 129–32, 135–6, 140, 142, 162; flaventior, 16, 128, 130, 132, 142; fraudans, 162; fuliginosa, 38; fuliginosa var. laetevirens, 38; fuliginosa var. laetevirens fo. eulaetevirens, 38; furfuracea, 16; furfuracea var. macrocarpa, 16; hababiana, 16, herreana, 16, 38; huachucensis, 16; hypoleucites, 16; hypopsila, 38; hypotropa, 16; incurva, 162; infumata, 162; isidiata, 16, 38; kurokawae, 16; leucochlora, 16; lineola, 16; livida, 16; **lophogena** comb. nov., 67–9; loxodes, 38; mellissii, 67–8; mexicana, 16, 38; molliuscula, 16, 38; mordenii, 68; multispora, 38; novomexicana, 16; olivacea, 16, 38; olivacea var. multispora, 38; omphalodes, 162, 166; panniformis, 38; paulensis, 68; physodes, 16, 39; plittii, 16, 39; praesignis, 16; praesorediosa, 68; prolixa, 16; pseudosulcata, 392; psoromifera, 16; pubescens, 39; pulla, 16; pulvinata, 16; quercina, 16; reticulata, 16; revoluta, 16; rudecta, 1–5; rutidota, 16; saxatilis, 39, 163; saximontana, 16; separata, 163, 165; sorediata, 39; sphaerosporella, 39; stenophylla, 17, 39; stygia, 39, 163; subargentifera, 17, 39, 48, 53; subaurifera, 39; subolivacea, 17, 39, 48, 53, 128–32, 135–6, 140, 142; subramigera, 17; subrudecta, 4, 17, 130, 132, 143; substygia, 17; subtinctoria, 17; sulcata, 4, 17, 39, 47–8, 53, 128, 130–2, 135, 143, 163, 165, 330; sulcata var. sulcata fo. ensulcata, 39; taractica, 17, 39; tasmanica, 17; tinctina, 17; trichotera var. trichotera, 39; tubulosa, 39; tucsonensis, 17; ulophyllodes, 17, 47–8, 53; verruculifera, 39; vittata, 39; weberi, 17; wyomingica, 17
Parmeliella microphylla, 17; praetermissa, 39, 161; saubinetii, 392, 399
Parmeliopsis aleurites, 17, 67; ambigua, 17, 39, 130–1, 133, 135, 140, 143, 467–8; hyperopta, 17, 39, 130–1, 133, 135, 140, 143, 392, 467, 469; placorodia, 17
Parmosticta, 185
Patellaria glomerulosa, 53
Peccania arizonica, 17
Pellia epiphylla, 75, 460
Peltigera, 181; aphthosa, 161, 165, 392, 398*,

399, 401; aphthosa var. aphthosa, 39; aphthosa var. variolosa, 17, 39; canina, 17, 39, 353, 394; canina fo. sorediata, 161; canina var. albescens, 17, 39; canina var. rufescens, 17, 39, 161; canina var. rufescens fo. innovans, 39; canina var. spuria, 39; canina var. spuria fo. sorediata, 39; canina var. ulorrhiza, 39; collina, 17, 39, 394; degenii, 39; horizontalis, 39; leucophlebia, 39; limbata, 39; malacea, 17, 39, 161; membranacea, 17, 39; polydactyla, 17, 39; praetexta, 39; pulverulenta, 161; rufescens, 39, 354; scutata, 39; spuria, 39; venosa, 17, 40

Peltula bolanderi, 17; clavata, 17; euploca, 17; obscurans var. deserticola, 17; obscurans var. hassei, 17; obscurans var. obscurans, 17; omphaliza, 17; placodizans, 17; polyspora, 17; richardsii, 17; zahlbruckneri, 17

Pennsylvania, mosses of, 458

Penzigiella, 423–5, 426*, 427, 429; cordata, 423–5, 427–9; robusta, 423; subcordata, *427*, 429

Pertusaria amara, 40; arborea, 40; bryontha, 162; coriacea, 162, 166; dactylina, 162; flavicunda, 17; glomerata, 162; granulata, 17; hymenea, 17; multipuncta, 17, 40; octomela, 162; panyrga, 162, 165; protuberans, 40; pustulata, 40; sommerfeltii, 18; subambigens, 392, 399; subobducens, 162, 165; trochisea, 162; wulfenii, 18; wulfenioides, 18

Peru, **Polytrichastrum** from, 201

Phaeoceros laevis subsp. carolinianus, 361

Phaeographis eulecta, 367

Phascum sp., 213

Philocrya aspersa, 357, 451

Philonotus fontana, 486

Physcia, 50, 54; adscendens, 18, 40, 48, 53, 130, 132, 135–6, 140, 143; aipolia, 18, 48, 53, 130–1, 133, 140, 143; aipolia fo. aipolia, 18, 40; aipolia fo. anthelina, 18; alba var. obsessa, 18; albicans, 18; albinea, 18; biziana, 18, 40; caesia, 18, 48, 53–4, 56, 163; callosa, 18; cernohorskyi, 54; ciliata, 18, 130, 133, 143; ciliata fo. ciliata, 18; ciliata fo. erythrocardia, 18; ciliata fo. fibrillosa, 18; convexa, 18; constipata, 18, 163; dubia, 40; endococcinea, 18, 48, 54; grisea, 18, 130, 132, 143; halei, 18; hirsuta, 44, 47–8, 50, 54, 56; hispida, 40; intermedia, 18, 163; melops, 18; millegrana, 4; muscigena, 18, 40, 163, 166; muscigena fo. isidiata, 40; muscigena fo. muscigena, 18, 40; muscigena fo. squarrosa, 163; nigricans, 40; orbicularis, 18, 40, 48, 54; orbicularis fo. albociliata, 54; orbicularis fo. orbicularis, 18; phaea, 18, 40; pulverulena, 18, 40; pulverulena fo. argyphaea, 40; pulverulena fo. pulverulena, 18; pulverulena fo. superfusa, 40; pulverulena fo. venusta, 40; sciastra, 18, 163; setosa, 18, 40; setosa fo. setosa, 18; setosa fo. virella, 18; stellaris, 4, 18, 40, 47–8, 54, 130, 135, 140, 143; stellaris fo. stellaris, 18; subtilis, 18; teretiuscula, 18, 40; tribacia, 48, 50, 54; wainioi, 18

Physciopsis syncolla, 48, 50, 54

Physconia sp., 48, 54; detersa, 40, 54; farrea, 54; grisea, 40; muscigena, 40; pulverulenta, 40

Physcomitrium, 438

Pike, L. H., W. C. Denison, D. M. Tracy, M. A. Sherwood and F. M. Rhoades: Floristic survey of epiphytic lichens and bryophytes growing on old-growth conifers in western Oregon, 389

Pilophoron acicularis, 40

Pilopogon longirostratus, 213; macrocarpus, 213

Pilotrichum wichurae, 427

Placodium amabile, 18; cinnabarinum, 18; elegans, 18; elegans var. brachylobum, 18; lobulatum, 18; murorum, 18

Placopsis gelida, 40, 162, 165

Placynthium aspratile, 161; microphyllizum, 18; nigrum, 18, 161; nigrum var. nigrum, 18; stenophyllum var. isidiatum, 18; subradiatum fo. coerulescens, 19

Plagiobryum zierii, 484–5

Plagiochila, 321; abdita, 205; abietina, 361; acanthocaulis, 205; alta, 362; asplenioides, 211, 306, 321, 325; major, 321; rhizophora, 321; satoi, 321; semidecurrens, 321; terebrans, 362

Plagiomnium cuspidatum, 486; medium, 486; venustum, 394

Plagiothecium cavifolium, 486; denticulatum, 486; laetum, 486; muellerianum, 94; turfaceum, 94

Platismatia glauca, 40, 330, 392, 398–400; herrei, 40, 392, 399; stenophylla, 40, 392, 400

Pleuridium, 152; sp., 213; andinum, 213; mexicanum, 213; nervosum, 158; palustre, 461

Pleuroclada, 322, 324; albescens, 324, 325

Pleuropus sericeus, 89

Pleurozia caledonica, 362

Pleurozium, 337–8; schreberi, 176, 215, 337, 484, 486

Pogonatum, 480; alienum, 480, 481*, 483; alpiniforme, *201*, 203; andinum, 212; arcuatum, 212; barnesii, *203*; bescherellei, 201, *203*; comosum, *201–4**, 483; cylin-

dricum, 202*–3; dentatum, 487; hirtellum, *203*; inclinans, 480; liebmannianum, 480, 482*–3; microstomum, 204; tenellum, *201*
Pohlia cruda, 486; flexuosa, 214; longicolla, 484, 486; nutans, 456–7, 486; wahlenbergii, 86
Polyblastia bryophila, 163; gelatinosa, 163; hyperborea, 163; integrascens, 163; theleodes, 163
Polyblastiopsis rupestris, 19
Polychidium muscicola, 40
Polyploidy, in Barbilophozia, 343
Polytrichaceae, anatomical studies, 113; neotropical, 201, 480
Polytrichadelphus ericoides, 213; longisetus, 213
Polytrichastrum, 480; alpiniforme, *201*; alpinum, 202, 487; alpinum var. septentrionale, 202; formosum, 483; papillatum, 201–2; **tenellum** comb. nov., 201–3*
Polytrichum, 113, 419, 480; alpestre, 415, 418–9; alpiniforme, *201*; alpinum, 487; comosum, *203*; commune, 57–62, 414, 416; commune var. commune, 487; commune var. maximoviczii, 484, 487; commune var. nigrescens, 487; formosum, 344; hyperboreum, 357; intermedium, *201*; juniperinum, 85, 343, 416, 487; piliferum, 57, 416, 461, 487; sexangulare, 487; strictum, 487; tenellum, *201*; yukonense, 358; spp. 288, 484
Porella, 324; cordaeana, 305, 324–6; navicularis, 324–6, 391, 394, 399–400; platyphylla, 324–5; platyphylloidea, 324–5; roellii, 324–6
Porothamnium expansum, 215
Porotrichum cobanense, 215; insularum, 215; wichurae, *427*–8
Pottia, 280
Pottiaceae, 247
Pottioideae, 282*
Prabhakar Naidu, C. S. *See* Roa, K. R., C. S. Prabhakar Naidu and N. Mala Kondaiah
Preissia quadrata, 222
Prionodon dichotomus, 214; fusco-lutescens, 214
Proctor, V. W. *See* Lipkin, Y. and V. W. Proctor
Protoblastenia rupestris, 18, 163
Protonema, of Bartramia pomiformis, 146
Pseudevernia furfuracea, 67; intensa, 19
Pseudobryum cinclidioides, 357, 461, 486
Pseudocrossidium, 279–280; apiculatum, 279; chilense, 279; elatum, 278
Pseudocyphellaria, 181; anomala, 40, 391, 393, 400; anthraspis, 40, 391, 394; crocata, 183, 393, 399; quercifolia, 183–6; rainierensis, 391, 394; sulphurea, 184
Pseudoditrichaceae, 489
Pseudoditrichum mirabile, 489–90
Pseudoscleropodium purum, 73–4
Pseudosymplepharis schimperiana, 213
Psilopilum, 447; cavifolium, 357
Psora crenata, 19; crenulata, 19; decipiens, 19; globifera, 19, 40
Psoroma hypnorum, 19, 40, 161, 165–6, 393, 401
Psorotichia schaereri, 19
Pterigynandrum filiforme, 486
Pterobryaceae, 423, 425; subfamily Trachylomoideae, 425
Pterobryon cordatum, *427*, *428*, 429
Pterygoneurum, 279–81; roseae, *253*
Ptilidium californicum, 307, 326; ciliare, 307–8, 325; cristacastrensis, 484, 486; pulcherrimum, 305, 307–8, 325, 460
Ptychomitrium gardneri, 371; serratum, 370–1
Pursell, R. A.: Distributional records of rare or infrequently collected species of mosses in Pennsylvania II, 93; and W. J. Hoe: Fissidens nothotaxifolius (Bryopsida: Fissidentaceae). A new species from the Hawaiian Islands, 476
Pylaisiella selwynii, 86
Pyramidula tetragona, 85
Pyrenodesmia cerina, 40
Pyrenopsis pulvinata, 19, 161; schaereri, 19
Pyrenula neglecta subsp. occidentalis, 367
Pyxine pringlei, 19

Racodium rupestre, 163
Radula, 324; auriculata, 324; bolanderi, 324, 394, 399–400; complanata, 211, 305, 324–5; obtusiloba, 324
Ramalina calicaris, 19; ecklonii, 19; farinacea, 19, 40; fastigiata, 19; fraxinea, 19; inflata, 40; intermedia, 19; pollinaria, 19; pollinariella, 40; roesleri, 40; sinensis, 19; yemensis, 19
Ramaut, J. L. and Mme M. Corvisier: Les acides lichéniques chez Stereocaulon virgatum, 365
Rao, K. R., C. S. Prabhakar Naidu and N. Mala Kondaiah: Chemical constituents of Riccia plana, 216
Reboulia hemisphaerica, 370, 460
Redon, J. (review) A. Hennsen and H. M. Jahns. Lichenes. Eine Einführung in die Flechtenkunde, 239
Reese, W. D.: Calymperes rubiginosum (Mitt.) Reese, com. nov., 92
Réunion, Anthocerotae and hepatics from, 361
Reviews: Hale, Jr., Mason E., The Biology

of Lichens, 386; Hawksworth, D. L., Mycologist's Handbook, An Introduction to the Principles of Taxonomy and Nomenclature in the Fungi and Lichens, 238; Henssen, A. and H. M. Jahns, Lichenes. Eine Einführung in die Flechtenkunde, 239; Instituton Botanicum, Acadamia Sinica (Editor) Iconographa Cormophytorum v. 1,2,3, 108; Kashyap, S. R., Liverworts of the Western Himalayas and the Panjab Plain, 239; Moore, P. D. and D. A. Bellamy, Peatlands, 386; Richardson, D. H. S., The Vanishing Lichens, Their History, Biology and Importance, 237; Universidad de Los Andes, Memoria del II Congreso Venezolano de Botánica (celebrado en Mérida entre junio 26 y julio 1 de 1972), 109

Rhabdoweisia crenulata, 359; crispata, 93; fugax, 221; perpusillum, 221

Rhacomitrium aciculare, 461; canescens, 485; ericoides, 484, 485; fasciculare, 485; heterostichum, 485; lanuginosum, 358, 484–5

Rhizocarpon badioatrum, 19; bisporum, 40; concentricum, 40; cookeanum, 40; copelandii, 162, 165; crystalligenum, 162; disporum, 19, 40, 162; eupetraeum, 40; geographicum, 19, 41, 162, 165; geographicum var. atrovirens, 41; grande, 19, 41; intermediellum, 19; jemtlandicum, 162; lecanorinum, 41; macrosporum, 19, 41; penichrum, 41; petraeum, 41; petraeum var. grande, 41; polycarpon, 19, 162; riparium, 19; rittokense, 162; sublucidum, 19

Rhoades, F. M. *See* Pike, L. H. et al.

Rhynchostegiopsis costaricensis, 469

Rhytidiadelphus, 484; triquetus, 187–8, 192, 394

Rhytidium, 337; rugosum, 337–8, 358–9, 486

Riccardia obtusa, 362; pinguis, 200, 211; **valida** comb. nov., 361–2; viridissima, 362

Riccia beyrichiana, 201; plana, 217–8

Riella subg. trabutiella, 25; affinis, 26; cossoniana, 25–30*; cossoniana var. echinata, 25, 27–30; helicophylla, 25–28*, 30; paulsenii, 25–29*; vishwanathai, 26

Riellaceae, 25

Rinodina sp. n. 49, 54–5; archaea, 47, 48, 50, 54, 394; coloradiana, 41; confragosa, 19; corticola, 19; darrovii, 19; euryspora, 19; exigua, 19, 48, 54, 130, 132, 143; finkii, 41; hallii, 41; laevigata, 19; marysvillensis, 41; milvina, 19, 163; mniaraea, 19; nigra, 41; nimbosa, 163; novomexicana, 19; occidentalis, 163; oreina, 19; pachysperma, 49, 55, 124, 129–30, 133, 143; pyrina, 47, 49–50, 55; roscida, 163, 166; sophodes, 19; thysanota, 19; turfacea, 163

Rozea bourgaenana, 215

Saelania glaucescens, 485

Sarcogyne clavus, 19; novomexicana, 19; pruinosa, 19; regularis, 20

Sayre, G.: Illustrations of the lost hepatics of the Wilkes Expedition, 204

Scapania, 317; americana, 318, 326; apiculata, 212, 305, 318, 325; arnellii, 318; bolanderi, 318, 326, 393, 397; carinthiaca, 318; curta, 305, 318, 325; cuspiduligera, 200, 318, 325; degenii var. dubia, 318; fissiloba, 205; glaucocephala, 318, 326; gymnostomophila, 318, 325; irrigua, 200, 318–9, 325; mucronata, 200, 318, 326; nemorosa, 318, 325, 459–60; paludicola, 318, 325; paludosa, 305, 319, 325; saxicola, 318–9, 326; scandica, 318; subalpina, 305, 318–9, 325; subalpina var. haynesiae, 318; uliginosa, 318; umbrosa, 305, 318–9, 325; undulata, 305–6, 318–9, 325, 456; undulata var. oakesii, 318

Scheirer, D. C.: Anatomical studies in the Polytrichaceae II. Histochemical observations on thickened lateral walls of hydroids of Dendroligotrichum, 113

Schistochila aligera, 362; caledonica, 362; stratosa, 362

Schroeder, G. L. *See* Schroeder, N. E., G. L. Schroeder and D. E. Anderegg

Schroeder, N. E., G. L. Schroeder and D. E. Anderegg: Catalog of the lichens of Idaho, 32

Sciaromium lescurii, 94

Scorpidium scorpioides, 176, 358, 486; turgescens, 176

Seligeria calcarea, 364; doniana, 364

SEM, on Riella spores, 28; of Alectoria, 78; on Sphaerocarpos donnellii spores, 81–2; of Thelia hirtella, 112; of Oligotrichum falcatum, 449; of Cyclodictyon richardsii, 470

Sematophyllum carolinianum, 95; demissum, 95, 461

Sendtnera quadrilaciniata, 205

Sesquiterpenoids, in Lophocolea heterophylla, 75

Seychelles, Anthocerotae and hepatics from, 361

Sharp, A. J. *See* Bowers, F. D. and A. J. Sharp; McNeilus, V. E. and A. J. Sharp

Sherwood, M. A. *See* Pike, L. H. et al.

Showman, R. E.: Lichens as indicators of air quality around a coal-fired power generating plant, 1

Sigal, L. L. and D. R. Toren: A new

Haematomma from California, 70; new distribution records of lichens in California II, 467
SMITH, D. K.: Three interesting and unusual mosses discovered in east Tennessee, 370
SMITH, G. L.: Neotropical Polytrichaceae I, II, 201; *See* STEERE, W. C. and G. L. SMITH; Neotropical Polytrichaceae III, 480
SNIDER, J. A.: Three new species of Archidium from Africa, 152
Solenostoma, 315; atrovirens, 316, 325; cordifolium, 305, 316–7, 325; crenuliformis, 316; gracillimum, 316, 325; hyalinum, 316, 325; oblongifolium, 316; obovatum, 316, 325; pumilum, 305, 316, 325; pumilum ssp. polaris, 316, 325; rubrum, 316, 326; sphaerocarpum, 305, 316, 325; triste, 316, 325
Solorina bispora, 161; crocea, 41, 181, 350; octospora, 161; saccata, 161; spongiosa, 161
South Dakota, hepatics of, 211, 304
SPESSARD, L. L.: Five mosses new to Illinois, 86
Sphaerocarpales, 25
Sphaerocarpos cristatus, 80; donnellii, 80–4*
Sphaerophorus globosus, 41, 163, 165, 393, 397, 398*–400
Sphagnum, 114, 321, 455–6, 458–9; 463–5, 489; spp., 57, 465; acutifolium, 463; angermanicum, 66; angustifolium, 65, 168, 175–7; aongstroemii, 357, 465*–7; balticum, 358, 463–4; capillaceum, 461, 485; centrale, 66; compactum, 357, 464, 485; cuspidatum, 65, 212, 461; cyclophyllum, 64; fallax, 65; fimbriatum, 65, 357, 455–7, 461; flaviomans, 66; flexuosum, 65; fuscum, 66, 464; girgensohnii, 65, 461, 485; imbricatum, 66, 357; jensenii, 65, 176; lenense, 358, 463–4; lescurii, 64; lindbergii, 65, 357, 465–6; macrophyllum, 65; magellanicum, 66, 168, 176–7, 461; majus, 65; molle, 64, 66; nemoreum, 65, 464; obtusum, 65, 357, 463; palustre, 461; papillosum, 66; perfoliatum, 463; pulchricoma, 65; pulchrum, 65; quinquefarium, 65; riparium, 65, 358; rubellum, 65, 464; russowii, 65; sparsum, 218, 221; squarrosum, 65, 456–7, 485; strictum, 64; subfulvum, 66, 358; subnitens, 66; subsecundum, 176, 358; subtile, 65; tabulare, 64; tenellum, 65, 218, 221; tenerum, 64–5, 218, 221; tenuifolium, 65; teres, 65, 357, 455–7, 485; trinitense, 65; vasculosum, 357; waghornei, 65; warnstorfii, 65, 168, 176–7; wulfianum, 65, 357
Splachnum turbinatum, 214
Spilonema revertens, 161
Sporastatia testudinea, 20, 162
Spores: of Sphaerocarpos donnellii, 80*; of Funaria hygrometrica, 194
Sporophyte, of Oligotrichum falcatum, 447
Squarmarina lentigera, 20
Staurothele catalepta, 20; clopima, 20; fissa, 20
STEERE, W. C. (eulogy) K. A. Holmen, 240; Occurrence of Entodon cladorrhizans in Alaska, with notes on geographical distribution of E. concinnus, 334; A new exsiccata from the northernmost parts of North America: Steere and Holmen, Bryophyta Arctica Exsiccata, Fascicle 1, Numbers 1–50, 356; Lophozia incisa: a curious reaction to desiccation, 368; and G. L. SMITH: The sporophyte of Oligotrichum falcatum, 447; and Z. IWATSUKI: Validation of Pseudoditrichaceae Steere et Iwatsuki fam. nov., 489
Stenocybe major, 393
Stereocaulon albicans, 20; alpinum, 162, 165; botryosum, 162, 165; denudatum, 365; paschale, 165; rivulorum, 162, 165; tomentosum, 41, 366; virgatum, 365–6
Sticta fuliginosa, 20, 393; weigelii, 20, 350–5, 393
Stokesiella oregana, 391, 393; praelonga, 215
Strangospora **microhaema** comb. nov., 44, 49–50, 55; **ochrophora** comb. nov., 44, 49–50, 55
Streptopogon erythrodontus, 214; matudianus, 221
SWINSCOW, T. D. V. *See* KROG, H. and T. D. V. SWINSCOW
Symphogyna crassifrons, 205
Synechoblastus coccophorus, 20; pycnocarpus, 20; texanus, 20
Syrrhopodon rubiginosus, 92

TAKEO, A.: Plasmolysis of protonema in Bartramia pomiformis, 146
Tayloria jamesonii, 214, 218, 221; moritziana, 218, 221; serrata, 363–4
Taxiphyllum deplanatum, 218, 221; machrisianum, 215
Taxonomic revisions: of Aloina, Aloinella and Crossidium, 245; of Penzigiella, 423
Taxonomic studies: Notes on Riella, subgenus Trabutiella, 25; three new species of Archidium, 152*; of Tomenthypnum falcifolium, 168; of Lecanora Sect. Petrasterion, 206; of Pseudoditrichaceae, 489
Teloschistes candlearius var. laciniosus, 41; modestus, 20; polycarpus, 20; ramulosus, 20
Temperature, effects on N-fixation in lichens, 350

Tennessee, Preissia quadrata from, 222; three mosses of, 370
Tetracymbaliella **comptonii** comb. nov., 361–2; ratiana, 362
Tetradontium, 364; brownianaum, 364
Tetraphis pellucida, 487
Tetraplodon angustatus, 364, 485; mnioides, 357, 485
Thamnolia vermicularis, 163, 165
Thelia hirtella, 112
THOMAS, R. J.: Sesquiterpenoids in the sporophyte of Lophocolea heterophylla, 75
THOMSON, J. W. (review) Hawksworth, D. L. Mycologist's Handbook. An Introduction to the Principles of Taxonomy and Nomenclature in the Fungi and Lichens, 238; *See* BARRETT, P. E. and J. W. THOMSON
Thrombium epigaeum, 41
Thuidium delicatulum, 456; peruvianum, 215; recognitum, 456
Thyrea nigritella, 367; pulvinata, 20
Thyridium rubiginosum, *92*
Timmia austriaca, 363–4, 484, 486
Tomenthypnum, 91; falcifolium, 168–77*; nitens, 168–9, 171–77*, 358; nitens var. falcifolium, 172; nitens var. involutum, 169
Toninia caeruleonigricans, 20; candida, 20; cinereovirens, 20; lobulata, 162; ruginosa, 20; tristis, 20, 467–9
TOREN, D. R. *See* SIGAL, L. L. and D. R. TOREN
Tortella fragilis, 364; humilis, 370
Tortula, 279–83, 285; aloides, *255*, 289; ambigua, *257*, 289; bifrons, *251*; brevirostris, *258*, 289; brevirostris var. elongata, *259*; calceolifolia, *261*; chloronotos, 274, *276*; crassinervia, *274*; cucullifera, *269*; duthieana, *261–2*; enervis, *261*; ericaefolia, 257; geheebii, *271*; goudotii, 214; membranifolia, *276*; muralis, 85; rigida, *255*, *261*; rigida ssp. aloides, *255*; rigida var. mucronulata, *262*; rigida var. obtusa, 262; rigida var. pilifera, *251*; rigida var. rigida, 262; ruraliformis, 62; ruralis, 370, 484–5; squamifera, *276*; squamifera var. grisea, *277*; squamifera var. pottioidea, *277*
TOWNSEND, C. C.: Pseudoscleropodium purum on two more tropical islands, 73
TRACY, D. M. *See* PIKE, L. H. et al.
Trachybryum megaptilum, 168
Trachyloma, 427
Trachylomoideae, 427
Trachypodopsis serrulata var. crispata, 221
Trebouxia, 472
Trentepohlia, 472
Trichostomum, 246; aloides, *255*; bartramii, 360; cylindricum, 94; tenuirostre, 94, 370
Tristichium mirabile, 218, 221
Tritomaria, 309, 314; exsecta, 212, 305, 325; exsectiformis, 305, 326; polita, 314, 325; quinquedentata, 314, 325; scitula, 212, 314, 325
TUCKER, S. C. and D. T. KOWALSKI: New state records of lichens from northern California, 366
TÜRK, R. and V. WIRTH: Über die SO_2-Empfindlichkeit einiger Moose, 187
Ulota crispa, 86; curvifolia, 357, 486
Ultrastructure, of Chiodecton sanguineum, 471
Umbilicaria angulata, 41; arctica, 162, 165; cinereorufescens, 20; decussata, 20, 41; deusta, 20, 41, 467–8; dillenii, 41; havaasii, 162; hirsuta, 20; hyperborea, 20, 41, 162; krascheninnikovii, 20, 41; papulosa, 20, 131–2, 143; phaea, 20, 41; phaea var. coccinea, 41; polyphylla, 41; polyrrhiza, 468; proboscidea, 20, 41, 162, 165; torrefacta, 20, 41; vellea, 20, 41, 162; virginis, 20, 41, 467–8
Urceolaria scruposa, 41; scruposa var. bryophila, 20
Usnea sp., 393, 400; aculeata, 20; arizonica, 20; cavernosa, 20, 131, 133, 143; ceratina, 20; comosa, 20; dasypoga, 20, 41, 330; erinacea, 20; florida, 20; florida fo. strigosa, 20; fulvoreagens, 21, 49, 55; glabrata, 41; glabrescens, 41; hirta, 21, 47, 49, 55, 128, 131–2, 135, 143; laricina, 41; mutabilis, 21; sorediifera, 21, 128–9, 131–2, 135–6, 143; subfloridana, 21; trichodea, 21; variolosa, 21
Utah, **Lecanora christoi** from, 206
Venezuela, additions to moss flora of, 212
Verrucaria arctica, 163; devergens, 163, deversa, 163, 166; fuscella, 21; margacea, 47, 49, 50, 55; muralis, 21; nigrescens, 21, 163; rupestris, 21; viridula, 21
Vestergrenopsis isidata, 161, 165
VITT, D. H. and C. D. HAMILTON: Taxonomic status of Tomenthypnum falcifolium, 168; *See* ANDRUS, R. E. and D. H. VITT; and R. E. ANDRUS, Sphagnum aongstroemii in North America, 463
WATSON, M. A.: Annual periodicity of incremental growth in the moss Polytrichum commune, 414
WEBER, W. A.: Two new species of Lecanora, Section Petrasterion, with a key to North American speices, 206
WEIN, R. W. and J. E. SPEER: Lichen biomass in Acadian and boreal forests of Cape Breton Island, Nova Scotia, 328

Weissia controversa, 363–4, 461
WIRTH, V. *See* TÜRK, R. and V. WIRTH
Wyoming, hepatics of, 304

Xanthoria candelaria, 21, 41, 163; elegans, 21, 41, 49, 55, 163, 165–6; fallax, 21, 41, 47, 49, 54, 56, 131–2, 143, 367; modesta, 21; polycarpa, 21, 41, 49, 54, 56, 131, 133, 143; substellaris, 21; sorediata, 21

Zygodon goudotii, 214; setulosus, 214

Volume 78, Number 3 (pages 245–388), was issued 4 December 1975.